


Kit Catalog number: 96-0293

Active Ingredient: 2-{2-Chloro-6-hydroxy-5-[2-methylquinolin-8-ylamino)methyl]-3-oxo-3H-xanthen-9-yl}benzoic acid (**FL**)
Catalog number - 07-0293



Introduction: The copper complex of **FL** is a novel, cell-permeable system that allows direct imaging of nitric oxide produced in living cells through fluorescence turn on. A solution of the copper (II) complex of **FL** can be readily prepared using this kit. For additional information, consult the following references:

1. *Nature Chemical Biology*, **2006**, 2, 375.
2. *Nature Protocols*, **2007**, 2, 408.
3. *J. Am. Chem. Soc.*, **2006**, 128, 14364.

Contents: Ligand **FL**: 5 x 0.5mg
Dimethylsulfoxide (ACS spectrophotometric grade): 5 x 1.0ml
Copper (II) chloride (1.0 mM solution in water): 5 x 1.0ml

MSDS: The Material Safety Data Sheets for the three products contained in this kit can be downloaded from the Strem Chemicals Web Site at www.strem.com. Locate the MSDS using the following catalog numbers:
FL: 07-0293
Dimethylsulfoxide (ACS spectrophotometric grade): 97-4940
Copper (II) chloride as a 1.0 mM solution in water: 97-3060

Storage conditions: The kit should be stored at -20°C and protected from light.

Preparation of the active copper complex of FL:

Step I

Allow the kit to warm to room temperature. Add 931 microliters of DMSO to a 0.5mg vial of **FL** (resulting concentration - 1.0 mM). The **FL** is readily soluble in the DMSO. The solution can be partitioned into aliquots of 40-300 µL as required. These solutions must be stored in the freezer at <-20°C. The DMSO solution of **FL** is stable for three months at -80°C. It is advisable to check the extinction coefficient of the solution before preparing the copper complex. (log ϵ (504 nm) = 4.6)

Step II

A Cu**FL** solution should be freshly prepared by adding 1:1 **FL** solution (1.0 mM) to the copper (II) solution (1.0 mM) at room temperature.

Note: The prepared DMSO/water stock solution of Cu**FL** solution [log ϵ (499 nm) = 4.6] can be kept at room temperature, but should be protected from light. The solution can be diluted with media to provide the concentration required for cell sensing experiments. When the extinction coefficient of the red solution of Cu**FL** diminishes by 20% of the original value, the solution should be discarded. Do not use the solution after 1 hour, and do not freeze the solution.

Visit www.strem.com for new product information and searchable catalog.

Strem Chemicals, Inc.
7 Mulliken Way
Newburyport, MA 01950-4098
U.S.A.
Tel.: (978) 499-1600
Fax: (978) 465-3104
Email: info@strem.com

Strem Chemicals, Inc.
15, rue de l'Atome
Zone Industrielle
67800 BISCHHEIM France
Tel.: (33) 03 88 62 52 60
Fax: (33) 03 88 62 26 81
Email: info.europe@strem.com

Strem Chemicals, Inc.
Postfach 1215
77672 KEHL
Germany
Telefon: 0 78 51/ 7 58 79
Email: info.europe@strem.com

Strem Chemicals UK, Ltd.
41 Hills Road
Cambridge
England CB2 1NT
Tel.: 0845 643 7263
Fax: 01223 368021
Email: enquiries@strem.co.uk